

# LA-UR-20-24201

Approved for public release; distribution is unlimited.

Title: Trinity Observations

Author(s): Carr, Alan Brady

Intended for: Material to be used as background during the Trinity anniversary.

Issued: 2020-06-09



### **Trinity Observations**

Various historical documents which record the observations of various scientists who observed the Trinity Test in southern New Mexico. The documents are contemporaneous with the test.

## To Be Completed by WRS-SIS staff, Collection Administrator, or CMC:

Note: This form is intended to be used for hand-carry purposes only. This step can be avoided if the document(s) are digitized and stored in an approved repository.

	mile) and institutes		an approved topository.			
Return to: Daniel Alcazar		Pho	one: 606-1563			
	Toro mono					
Date	2/6/2020					
Document Unique	A-84-019 (16/5)					
Identifier	10					
Page or Image Count	12					
Requestor Name and Z#	Alan Carr Z# 195680					
	=======================================					
Requested Return Date	5/28/2020					
<ul> <li>Other Transfer/Collection No.:</li> <li>Notify SAFE-IP coordinator of electronic red file transfer through yellow email.</li> <li>Date: 5/27/2020</li> <li>To Be Completed by Classification Group:</li> </ul>						
Date Received:	Date Received:		Date Released:			
Classification Category an	d Level:					
СП	S		RD 🗀			
FRD	NSI 🗆		Sigma/Other			
Unclassified*	Caveat:					
Comments: Lovered RODE WT ENS.						
Document requires derivative classifier (DC)-verified unclassified stamping.						
Joop 52 July			MAYLOLO			
Classification Analyst Signatu	re	Date				

Release date: January 16, 2019

27 July 1945.



### Description of Trinity Test, as Observed From Coordinating Council Area

- 1. The area assigned to the Coordinating Council, from which I observed the Trinity Test Shot, was understood to be approximately 20 miles from point of detonation. The intervening landscape was quite flat, and a clear but distant view of the explosion was possible.
- 2. At the mement of detonation I was seated on the ground facing in the general direction of the test setup, with my eyes fixed on the ground immediately in front. This I did deliberately, in order to avoid the blinding flash which I expected at the instant of detonation.
- 3. My first impression was of sudden brilliant lighting of the surrounding landscape, accompanied by a momentary flash of heat. I remember a feeling of surprise as the illumination, initially quite brilliant, continued to increase for a brief interval. After raising the dark filter to protect my eyes, I looked in the direction of the light. Although the filter provided was designed to eliminate over 99% of the light, the intensity of illumination was such that there was a momentary sensation of blinding, similar to that following a close flash of lightning on a dark night.
- 4. Within perhaps two seconds after the initial flash, it became possible to distinguish details of the explosion without the use of the filter. A ball of light about three or four hundred yards in diameter, was clearly evident about a thousand feet above the ground. Beneath this ball there appeared to be a column of red flame about 150 or 200 yards in diameter. Flickering red reflections were distinctly seen on the clouds above the ball of light.
- 5. As the intensity of light diminished, a smokey, grayish-brown ball took shape. It was noted that a fringe of violet light surrounded this ball and that the ball was rising rapidly, with some increase in diameter. The column beneath the ball darkened until it appeared to be a dense black pillar under the grayish-brown ball. At about ten seconds after detonation, the top of the ball seemed to flatten, and the ball and column took on the shape of a vast mushroom.
- 6. Close to the ground and slowly spreading out on all sides from the point of detonation, was a white cloud which looked much like ground fog. This cloud, I assumed, was composed of dust raised by the blast wave. The maximum diameter of this cloud I estimated to be about two miles.

VERIFIED UNCLASSIFIED

LMR 7-17-79

GMS JUL 9 1981

JLASSIFIE

SPECIAL RE-REVIEW
FINAL DETERMINATION 1981
UNCLASSIFIED, DATE:

Lt. Taylor

= 2 =

27 July 1945.

7. The sound wave when it reached my position, was unexpectedly sharp. I would describe the sound as being similar to the crack of thunder following a near stroke of lightning. It was sharper and lacked the booming sound of a large gun firing. There was no perceptible earth shock felt at any time.

8. Ten minutes after the detonation the mushroom-like cloud was still quite distinct and rising rapidly, with very little tendency towards dissipating horizontally. About fifteen minutes after the explosion, the pillar under the cloud had faded. The cloud, now a light brownish color, began to fade. The cloud faded from view at about thirty minutes after the explosion.

RAL/hg

R. A. Larkin Captain, USN

Cas Ugg mheimen

UNCLASSIFIED

		Olyonation				
Clas	authority	of the	U. S. Atomic E	nergy Commissi	on,	
Per	(Person	authoris	ing change in	classification)	(Date)	
BZ	22	K.	of person mak	ing the change,	4-3-75 and date)	

UNCLASSIFIED

CLASSIFICATION CANCELLED PER DOC REVIEW JAN. 1973

U. S. ENGINEER OFFICE ROJECT "YAUG 2545

25 AUGUST 1945

SPECIAL RE-REVIEW FINAL DETERMINATION DATE :

Washington Liaison office WASHINGTON, D. C.

FROM: CLEAR CREEK, N. M.

VERIFIED UNCLASSIFIED

THE RADIOACTIVITY OF THE SOIL AT TRINITY WAS QUITE HIGH EUT WAS COMFINED ALMOST ENTIRELY WITHIN A RADIUS OF TWELVE HUNDRED YARDS PD PAREN FOR GENERAL L R GROVES FROM DR LOUIS HEMPELMANN PAREN AFTER TWENTY FOUR HOURS THE HIGHEST ESTIMATE OF RADIOACTIVITY IN THE CRATER BY MEMBERS OF THE TANK CREW WAS SIX FIVE ZERO R PER HOUR PD THIS FELL TO TWO R PER HOUR AT EIGHT ZERO ZERO YARDS AND TO ONE SLANT TEN R PER HOUR AT ONE TWO ZERO ZERO YARDS PD AFTER ONE WEEK THE ABOVE ACTIVITY HAD DECAYED BY A FACTOR OF FIFTEEN PD AFTER THIRTY DAYS THE HIGHEST RADIATION INTENSITY WAS FIFTEEN F PER HOUR PD THE AREA CONTAINING SIGNIFICANT ACTIVITY PAREN MORE THAN XXX ONE SLANT TEN R PER HOUR PAREN HAD SHRUNK TO A RADIUS OF FOUR ZERO ZERO YARDS PD THE RADIATION WAS DUE TO BOTH INDUCED RADIOACTIVITY OF THE SOIL RESULTING FROM THE ACTION OF NEUTRONS AND TO THE DEPOSITION OF FISSION PRODUCTS ON THE GROUND PD THE INDUCED RADIOACTIVITY WHICH CORRESPONDED TO ABOUT ONE SLANT TWO OF THE TOTAL AT ONE DAY CMA WAS CHIEFLY RADIOACTIVE SODIUM PD IT DECAYED VERY RAPIDLY PAREN HALF DASH LIFE FIFTEEN HOURS PAREN AND WAS INSIGNIFICANT AFTER ABOUT ONE WEEK PD THE FISSION PRODUCTS ARE DISAPPEARIVE MORE SLOWLY FOLLOWING ONE SLANT T LAW PD RADIOACTIVE MATERIALS DID NOT FALL FROM THE CLOUD OUTSIDE THE ABOVE AREA IN AMOUNTS WHICH WOULD BE CONSIDERED DANGER CUSASS

15 45

CV

LO C



GREAT HEIGHT OF DETONATION THE NEUTRON INDUCED ACTIVITY OF THE SOIL SHOULD BE INSIGNIFICANTA DEPENDING UPON METEOROLOGIC CONDITIONS CMA THE ACTIVITY DUE TO FISSION PRODUCTS MAY BE MUCH LESS OR MUCH GREATER THAN AT THE TRINITY TEST

Approval for Transmission:

Ru daylol



CLASSIFICATION CANCELLED PER DOC REVIEW JAN. 1973

U. S. ENGINE PROJECT

25 AUGUST 1945

WASHINGTON LIAISON OFFICE WASHINGTON, D. C.

FROM: CLEAR CREEK, N. M.

VERIFIED UNCLASSIFIED

THE RADIOACTIVITY OF THE CRATER REGION AT TRINITY WAS CONFINED TO A ROUGHLY CIRCULAR AREA OF ONE TWO ZERO ZERO YARDS RADIUS PD PAREN FOR GENERAL I R GROVES FROM LOUIS HEMPELMANN INFORMATION FOR COL WARREN PAREN THE HIGHEST RADIATION INTENSITY WAS ESTIMATED TO BE SIX FIVE ZERO R PER HOUR IN THE CRATER AND WAS MEASURE TO BE TWO R PER HOUR AT EIGHT HUNDRED YARDS AND ONE SLANT TEN R PER HOUR AT TWELVE HUNDRED YARDS PD THE ACTIVITY WAS SPOTTY BEING HIGHEST OVER THE GREEN GLAZED MATERIAL PD THE RADIATION FROM THE GROUND WAS DUE TO RADIOACTIVE : SODIUM AND TO FISSION PRODUCTS IN ABOUT EQUAL AMOUNTS AT TWENTY FOUR HOURS CMA THE SODIUM HAD DISAPPEARED AND THE RADIOACTIVE DECAY FOLLOWED THE ONE SLANT T LAW PD AT THIRTY DAYS CMA THE HIGHEST RADIATION INTENSITY WAS FITTEEN H PER HOUR AT THE PERIPHERY OF THE CRATER PD THE KADIUS OF THE AREA OF SIGNIFICANT RADIATION PAREN MORE THAN ONE SLANT TEN R PER HOUR PAREN HAD SHRUNK TO FOUR ZERO ZERO YARDS PD IN COMBAT USE CMA DUE TO THE HEIGHT OF DETONATION CMA THERE SHOULD BE LITTLE IF ANY NEUTRON INDUCED ACTIVITY OF THE SOIL PD THE ACTIVITY DUE TO FISSION PRODUCTS SHOULD BE GREATER OR LESS THAN THAT AT TRINITY DEPENDING UPON METEOROLOGIC

CONDITIONS CMA ESPECIALLY HUMIDITY

PER DOC REVIEW JAN. 1973

Approval for Transmission: Kin day los

00

CV

inis document consists of.....rage(s of \_\_\_\_\_Copies, Series Q

by Observations During the Explosion at Trinity on July 16, 1945 - E. Fermi

On the morning of the 16th of July, I was stationed at the Base Camp at Trinity in a position about ten miles from the site of the explosion.

The explosion took place at about 5:30 A.M. I had my face protected by a large board in which a piece of dark welding glass had been inserted. My first impression of the explosion was the very intense flash of light, and a sensation of heat on the parts of my body that were exposed. Although I did not look directly towards the object, I had the impression that suddenly the countryside became brighter than in full daylight. I subsequently looked in the direction of the explosion through the dark glass and could see something that looked like a conglomoration of flames that promptly started rising. After a few seconds the rising flames lost their brightness and appeared as a huge pillar of smoke with an expanded head like a gigantic mushroom that rose rapidly beyond the clouds probably to a height of the order of 30,000 feet. After reaching its full height, the smoke stayed stationary for a while before the wind started dispersing it.

About 40 seconds after the explosion the air blast reached me. I tried to estimate its strength by dropping from about six feet small pieces of paper before, during and after the passage of the blast wave. Since at the time, there was no wind I could observe very distinctly and actually measure the displacement of the pieces of paper that were in the process of falling while the blast was passing. The shift was about 22 meters, which, at the time, I estimated to correspond to the blast that would be produced by ten thousand tons of T.N.T.

classification) (Date) aking the change, and date PUBLICLY RELEASABLE

10

CV.

1

N

N

LANL Classification Group

RE-REVIEW

SPECIAL RE-REVIEW

This Document Contains FFICE MEMORANDUM

Copies.

July 20, 1945 DATE

TO:

FROM:

Joe Hoffman

John Blair

Max Kupferberg

Section I-6 Searchlight Detachment

Alex Nedzel

SUBJECT: Cloud Observations and Radiation Measurements

VERIFIED UNCLASSIFIED

The report includes data on the position of the cloud and on the radiation as measured by Searchlight Section I-S, situated 19.5 miles from position zero, 330 east of north. Azimuth and elevation angles are presented, as well as a description of the phenomena seen. The azimuth and elevation readings were combined with similar readings from Sl. Sec. 1-7, 1-2 and 1-3 to determine the course of the cloud. The information was radiced to I-l, where the actual course was plotted by Capt. Allen. Measurements were taken on azimuth, elevation, 3 and 7 radiation, starting at time, t = 0.

At time t = 0, all three observers were startled by the high intensity of the light from the explosion. They independently reported feeling the heat radiation on their faces. The last part of the ball of fire then rose above the hills between positions L-6 and Q. It was decided that it rose at least two diameters above the hills. It was at this point that the actual cloud became visible. It seemed to move directly along a vertical until the upper portion formed a dense white mushroom. The mushroom was trailed by a dusty red-brown streamer.

The radiation meters were closely watched from t = 0 until the cloud was well into the sky. They indicated no increase in radiation. It might be noted that the part of the ball of fire seen from this position had very nicely formed billows in it. Once the cloud was sighted, readings on elevation, azimuth and radiation were recorded periodically (See Table I).

The radiation was measured both on the ground and approximately three feet above the ground. The major part of the radiation measurements were done within a fifty-yard square. Some measurements were taken at a mile east and a mile CANCELLED position L-S and found to agree with the readings of position L-S and the readings of position L-S an PER DOCTREVIEW IAN. 1973 were a Landsverk and Wollan ( and V ) quartz-fiber electrometer and a Victoreen V

Form 25

1

N

117

N

To:

Joe Hoffman

本家

John Blair Max Kupferberg Alex Nedzel

SUBJECT:

FROM:

Cloud Observations and Radiation Measurements

FINAL DETERMINATION UNCLASSIFIED

L. M. Redman

JUL 17 79

survey meter.

The appearance of the cloud from the position I-C is of interest. It was divided into three parts before the time t = / 15 minutes a dense white mushroom cloud, a flat fairly long red dust cloud, and a reddish-brown column which seemed to come from position 0. At the time / 30 minutes, the high mushroom cloud had moved directly towards position L-E, and had taken on the shape of the North American part of the western hemisphere. The lower red-brown cloud and column took on the shape of a question mark, while the brown dust seemed to be still emanating from position O. This was verified by the three observers. There was still no indication of Y or /3 radiation. At the time t = / 55 minutes, the mushroom part of the cloud was directly overhead and spreading out rather than moving to the east, as was the case between the time when the mushroom was formed and t = # 55 minutes. The radiation present at t = 55 minutes was not enough to be recorded on the Landsverk and Wollan electrometer. From t = / 55 minutes until the cloud was obscured by a lower formation of clouds, it stayed at the zenith with respect to position L-S. Even when the cloud was obscured it was known that it did not move a great deal as a cloud, because the radioactive cloud was never seen to move from behind the lower formation of clouds. It seemed as though the cloud moved up overhead and was dispersed until its density was low enough to make it indistinguishable from other thin clouds. This doesn't mean that the material making up this cloud stayed over the region of position I-S. It spread over the whole area.

The radioactive cloud was hidden at t = 1 hour and classification CANCELLED and elevation readings could no longer be taken, but \( \sigma \) and \( \sigma \) radiperson readings were continued.



m

N

N

DATE

July 20, 1945 - Page 3

#### INTER-OFFICE MEMORANDUM

To:

Joe Hoffman

John Blair

Wax Kupferberg

Alex Nedzel

Cloud Observations and Radiation Measurements

Radioactive material started to descend upon the position L-6 between the time / 1 hour, 30 minutes and 2 hours, 0 minutes, thirty-five minutes to an hour after it reached the zenith. At 2 hours, 0 minutes the radiation had increased to .10 R/hr. Before # 1 hour, 30 minutes the radiation was less than 1 m R/hr. The 7 radiation measurement was made 1" above the ground level. The gamma radiation increased to 2.0 R/hr. at t = # 2 hours, 55 minutes. It then leveled off and continued at 2.0 R/hr. until time t = 3 hours, 45 minutes. The curve of R/hr. vs. time then shows a decay.

Radiation readings were taken both on the ground and three feet above it on Ys and /3s / Ts, respectively. Readings on Ys, taken three feet above the ground were 25% less than of readings taken 1" above the ground (See Table I: / 2 hours, 55 minutes). When the Y radiation on the ground was 2.0 R/hr., the /3 activity on the ground was above 5.0 R/hr. The / / Breadings on the ground remained greater than 5 R/hr. for the duration of the readings. At the beginning of the descension, the # /3 activity was not appreciably larger than the Y activity alone. At the time 4 hours, 45 minutes, the Victoreen 7 meter was loaned to the health group and all subsequent measurements were made with the L & W meter.

In addition to the data recorded at position I-2, Yradiation at Bingham, four miles west of Bingham, at White's store and at I-6 was recorded. The readings and the time at which they were taken are presented in Table I. John Magee reported 15.0 R/hr. Tradiation off the ground 3.5 miles east of L-8. This reading was checked and found to be correct. It was recorded at # 3 hours, 20 minutes.

As an interesting point it should be noted that at the time the average reading was .1 R/hr. there were small areas one foot square which provide the square which provi



0

N

10

CA



UNCLASSIFIE

DATE July 20, 1945 - Page 4

#### INTER-OFFICE MEMORANDUM

To:

Joe Hoffman

XXX

John Blair

FROM:

Max Kupferberg

Alex Nedzel

SUBJECT:

Cloud Observations and Radiation Measurements

The readings taken with the radiation meters have an accuracy of  $\angle$  10%. The accuracy of the measurements are probably typical for the type of meters used. These meters were obtained from R. Watts, who is undoubtedly familiar with the accuracy of their calibrations.

Copies to: Bacher

Hershfelder Hoffman Kupferberg Nedzel Oppenheimer Warren

MILASTEL

CLASSIFICATION CANCELLED PER DOC REVIEW JAN. 1978



N

In

N

N